



Agriculture
Canada



Agriculture
Canada

Canadian Agriculture Library
Bibliothèque canadienne de l'agriculture
Ottawa K1A 0C5

23



Research Branch
Technical Bulletin 1993-17E

Guide to the Wild Germplasm of Brassica and Allied Crops

Part IV

Centre for Land
and Biological Resources Research



Centre de recherches sur les
terres et les ressources biologiques

Canada

50.72
759
73-17

Cover illustration

The images represent the Research Branch's objective: to improve the long-term competitiveness of the Canadian agri-food sector through the development and transfer of new technologies.

Designed by Research Program Service.

Illustration de la couverture

Les dessins illustrent l'objectif de la Direction générale de la recherche : améliorer la compétitivité à long terme du secteur agro-alimentaire canadien grâce à la mise au point et au transfert de nouvelles technologies.

Conception par le Service aux programmes de recherches



Guide to the Wild Germplasm of Brassica and Allied Crops

Part IV Wild Species in the Tribe Brassiceae (Cruciferae) as Sources of Agronomic Traits

S.I. Warwick
Centre for Land and Biological Resources Research
Ottawa, Ontario

Technical Bulletin 1993-17E

Centre for Land and Biological Resources Research
Research Branch, Agriculture Canada

November 1993

Copies of this publication are available from:

Centre for Land and Biological Resources Research
Research Branch, Agriculture Canada
K.W. Neatby Bldg., C.E.F.
Ottawa, Ontario
K1A 0C6

Published by Cartographic Design and Reproduction Unit
Centre for Land and Biological Resources Research

© Minister of Supply and Services 1993
Cat. No. A54-8/1993-17E
ISBN 0-662-21117-0

ACKNOWLEDGEMENT

I wish to thank L.D. Black, Centre for Land and Biological Resources Research for her technical assistance in preparing this publication.



Digitized by the Internet Archive
in 2013

<http://archive.org/details/guidetowildgerm199317warw>

INTRODUCTION TO THE GUIDE:

The Cruciferae family, which contains about 3500 species and 350 genera, is one of the ten most economically important plant families (Rich 1991). The tribe Brassiceae is one of the 13-19 tribes which have been recognized within the family and is one of the few tribes believed to constitute a natural group (Hedge 1976, Al-Shehbaz 1984, 1985). It is the most important economically and the most distinctive (Gómez-Campo 1980, Al-Shehbaz 1985). It is distinguished on the basis of the presence of conduplicate cotyledons (i.e. the cotyledons are longitudinally folded around the radical) and/or two-segmented fruits (siliques) which contain seeds in one or both segments, and only simple hairs if present (Gómez-Campo 1980, Al-Shehbaz 1985).

Crop brassicas display enormous diversity and are used as a source of oil, vegetables, mustard condiments, and fodder. Those of particular importance in Canada are: *Brassica napus*, *B. rapa*, and *B. juncea* as sources of canola oil, and *B. oleracea* as cole-crops. The genera *Raphanus* and *Sinapis* are also of major importance, the former cultivated for its edible roots and the latter as a source of mustard condiments along with *B. nigra*. Several species have become naturalized weeds in Canada and the United States [eg. *Sinapis arvensis* (wild mustard), *Raphanus raphanistrum* (wild radish), and *B. rapa* (wild rape)], representing both a potential source of germplasm and agricultural problems. In other areas of the world *Crambe* is cultivated as an industrial oil, and the leaves of other genera (eg. *Eruca* and *Diplotaxis*) are eaten as salad greens.

An understanding of the genetic potential of wild relatives of the crop species of *Brassica* and allied genera (members of the Tribe Brassiceae) is critical for the establishment of long-term breeding programs of these crops. In addition, it is clear that many of the wild species in the tribe have potential value as new crops, as sources of industrial oils (*Crambe*, *Eruca*), condiments (*Sinapis alba*), and other diverse products. Wild relatives also possess a number of useful agronomic traits which could be incorporated into breeding programs, including: cytoplasmic and nuclear male sterility; resistance to disease and insect and nematode pests; intermediate C₃-C₄ photosynthetic activity; and tolerance of cold, salt and drought conditions.

The last comprehensive taxonomic treatment on the tribe was conducted by Schulz (1919, 1923, 1936). The tribe Brassiceae contains approximately 217 species and 51 genera, 25 of which are monotypic (Table below). Geographically, it is centered in the southwestern Mediterranean region, particularly Algeria, Morocco and Spain, where c. 41 genera are either endemic or exhibit maximum diversity. The tribal range extends eastward into India and Pakistan and southward into South Africa, with a poor representation in the New World (Hedge 1976, Gómez-Campo 1980, Al-Shehbaz 1985).

GENERA OF THE TRIBE BRASSICEAE (no. species in brackets)

Ammosperma (2)	Guiraoa (1)
Boleum (1)	Hemicrambe (2)
Brassica (35)	Henophyton (1)
Cakile (7)	Hirschfeldia (2)
Carrichtera (1)	Kremeriella (1)
Ceratocnemum (1)	Moricandia (9)
Chalcanthus (2)	Morisia (1)
Coincya (6)	Muricaria (1)
Conringia (6)	Otocarpus (1)
Cordylocarpus (1)	Physorrhynchus (2)
Crambe (26)	Pseuderucaria (2)
Crambella (1)	Pseudofortuynia (1)
Didesmus (2)	Psychine (1)
Diplotaxis (27)	Quezeliantha (1)
Dolichorhynchus (1)	Raffenaldia (2)
Douepia (1)	Raphanus (2)
Enarthrocarpus (5)	Rapistrum (2)
Eremophyton (1)	Rytidocarpus (1)
Eruca (3)	Savignya (1)
Erucaria (9)	Schouwia (1)
Erucastrum (19)	Sinapidendron (5)
Euzomodendron (1)	Sinapis (5)
Fezia (1)	Succowia (1)
Foleyola (1)	Trachystoma (3)
Fortuynia (2)	Vella (5)
	Zilla (1)

Within the tribe, Schulz (1919, 1923, 1936) also recognized, somewhat arbitrarily on the basis of morphological characters, seven subtribes: Brassicinae, Cakilinae, Moricandiinae, Raphaninae, Savignyinae, Vellinae, and Zillinae. Gómez-Campo (1980) has since proposed a reduction to six subtribes with the inclusion of the Savignyinae in the Vellinae. The Brassicinae and Moricandiinae both include genera with elongated siliquose dehiscent fruit, while the other subtribes include those with reduced or "nucamentaceous" fruits.

Generic boundaries in the tribe are still somewhat arbitrarily drawn, and the establishment of clear-cut intergeneric relationships requires clarification. Unlike many of the small genera, the species are generally very distinct throughout the family, with fruit characters being the most reliably used structures for the proper identification of genera and species. Taxonomic debate in the tribe has centred most particularly upon the number of and relationships between the subtribes and genera (Hedge 1976, Al-Shehbaz 1985).

The genus *Brassica* is one of ten core genera in the subtribe Brassicinae, which also includes *Coincya*, *Diplotaxis*, *Eruca*, *Erucastrum*, *Hirschfeldia*, *Raphanus*, *Sinapidendron*, *Sinapis*, and *Trachystoma*. The Brassicinae is defined primarily on the basis of elongated (siliquose) dehiscent fruits, presence of median nectaries, and usually seeded beaks. Although morphologically quite distinct from subtribes Cakilinae, Vellinae, and Zillinae, its separation from the Raphaninae and Moricandiinae is less clear. Current generic circumscriptions within the subtribe Brassicinae have also been considered to be highly artificial by many taxonomists, with generic delimitation based primarily on only one or two morphological traits.

Systematists are continuing to re-evaluate relationships within the tribe Brassiceae by way of morphological, cytological, hybridization, isozyme and molecular analyses (studies reviewed in Warwick and Black 1991, 1993). Such research has confirmed many proposed species relationships, but has also indicated new relationships between genera and species. In particular, these studies have identified new potential sources of germplasm for *Brassica* crops, indicating that the range of germplasm important to the genus is much greater than previously recognized.

The following Guide to the wild germplasm of *Brassica* and allied crops (Tribe Brassiceae, family Cruciferae) will be divided into five parts as indicated below:

- I. **Taxonomy and Genome Status** - [Complete list of genera and species in the tribe and their genomic status, containing cross references for commonly confused names];
by S.I. WARWICK
- II. **Chromosome Numbers**;
by S.I. WARWICK & J.K. ANDERSON
- III. **Interspecific and Intergeneric Hybridizations**;
by S.I. WARWICK & L.D. BLACK
- IV. **Wild Species as Sources of Agronomic Traits** - [List of potentially useful agronomic traits and possible wild germplasm sources in the tribe];
by S.I. WARWICK
- V. **Life History Data** [Summary of habitat and geographical distributions of all species indicated in part I.]
by S.I. WARWICK

The information provided in this guide is intended to be useful in providing direction for future genebank needs for these crops and for assisting biotechnologists and breeders wishing to utilize these genetic resources in their research programs.

PART IV. WILD SPECIES IN THE TRIBE BRASSICEAE (Cruciferae) AS SOURCES OF AGRONOMIC TRAITS

The following publication is the fourth part of a guide to the wild germplasm of *Brassica* and allied crops (Tribe Brassiceae, family Cruciferae).

The potential of wild members of the tribe Brassiceae as sources of agronomic traits will be reviewed. In addition to traditional breeding methods, interspecific and intergeneric transfer of genes governing qualitative and quantitative characters from wild allies to cultivated forms will be facilitated with various in vitro methods, such as somatic cell genetics and recombinant DNA techniques (Sjödin 1992).

Examples of genetic variability in potential agronomic traits of germplasms of *Brassica* and related genera will be presented under the following sections:

- 1. North American Germplasm**
 - A. North American Ethnobotany
 - B. Weed/Crop Genetic Exchange
 - C. Weed and Crop Species in Canada and U.S.A.
- 2. Morphological Characters**
 - A. Hairs on cotyledons, juvenile leaves, fruits
 - B. Leaf thickness/waxiness
 - C. Petal colour
 - D. Resistance to pod shattering
 - E. Yellow Seed
 - F. Growth form
- 3. Chemical Traits**
 - A. Fatty Acids
 - B. Glucosinolates (mustard oil glucosides)
 - C. Phenolic Choline Esters
- 4. Photosynthesis**
- 5. Cytoplasmic Male Sterility**
- 6. Soil Adaptation:**
 - A. Saline
 - B. Calcareous
- 7. Cold Tolerance**
- 8. Drought Tolerance**
- 9. Herbicide Resistance**
- 10. Disease Resistance**
- 11. Insect Resistance**
- 12. Nematode Resistance**
- 13. New Crops**

1. North American Germplasm:

The genus *Brassica* is considered a native of Eurasia and to have been introduced into the New World by European settlers around 400 years before present (B.P). Recent archaeological and ethnobotanical studies, reviewed in Jacobson et al. (1988), have provided evidence for a North American distribution of *Brassica*. *Brassica kaber* [= *Sinapis arvensis*] was found to have occurred in the northeastern United States as early as 8000 B.P. and was abundant and widespread by 2000 B.P.. These suggest the original existence of a semi-circumboreal distribution for *B. kaber* (and perhaps other *Brassica* spp.) which was maintained during the late-Quaternary with the range in North America changing during the glacial/interglacial cycles.

North American germplasm of the Tribe Brassiceae, whether naturalized introductions/crop escapes or native components, represents both a wild source of agronomic traits as well as agricultural weed problems.

Historically Native peoples of North America have used a number of "wild" *Brassica* spp. for both food and medicinal purposes:

A. North American Ethnobotany: (Arnason et al. 1981, Jacobson et al. 1988)

- | | |
|--|--|
| <i>Brassica</i> spp. | - young shoots cooked as greens by Iroquois & Malecite Indian Tribes. |
| <i>Brassica nigra</i>
(black mustard) | - seeds ground and used as snuff to cure head colds by the Meskwaki.
- leaves used to relieve toothaches and headaches by the Mohegans. |
| <i>Brassica hirta</i>
(white mustard) | - used to treat tuberculosis by the Malecite and Micmac |
| <i>Brassica napus</i>
(wild turnip) | - bark used to treat colds, cough, grippe and small pox by the Micmac
- used for chilbains by the Rappahannock |
| <i>Brassica oleracea</i>
(wild cabbage) | - used for headaches by the Rappahannock |
| <i>Brassica rapa</i> | - was used as medicine by the Bois Fort Chippewa |

B. Weed/Crop Genetic Exchange:

Weed/crop interchange among members of the same cytodeme is possible and in studies on the origin of variation in wild *Raphanus sativus* in California Panetsos & Baker (1968) suggested that the introgression of wild radish, *R. raphanistrum*, characters appears to have been a major factor in converting the former crop, *R. sativus*, into a highly successful weed. Similar possibilities exist with crops grown in an area with dense populations of naturalized or escaped members of the same species or cytodeme.

Of more recent concern is the risk assessment of gene transfer between crops and weeds (intercytodeme exchange) [See PART III of the GUIDE].

These include:

- Sinapis arvensis* with *Brassica nigra* and *B. juncea* (Bing et al. 1991);
- Hirschfeldia incana* (= *Brassica adpressa*) with *Brassica napus* (Lefol et al. 1991);
- Raphanus raphanistrum* and *Hirschfeldia incana* with *Brassica napus* (Kerlan et al. 1991).

C. Weed and crop species in Canada & United States:

Weed and crop species [Common name] in Canada:	Distribution
- <i>Brassica juncea</i> (L.) Czern & Coss. [Indian mustard]	NT-M NF NS PE NB PQ ON MB SK AB BC
- <i>Brassica napus</i> L. [rape, canola]	NT-M LB NF PE NS NB PQ ON MB SK AB BC
- <i>Brassica nigra</i> (L.) Koch [black mustard]	NF SM NS PE NB PQ ON SK BC
- <i>Brassica oleracea</i> L. [cabbage]	NF PE PQ ON
- <i>Brassica rapa</i> L. [bird rape] (= <i>Brassica campestris</i> L.)	NT-M YT LB NF NS PE NB PQ ON MB SK AB BC
- <i>Cakile</i> spp." [sea-rocket]	NF NS PE NB PQ ON BC
- <i>Conringia orientalis</i> (L.) Andrzejowski [hare's-ear mustard]	NF NS PE NB PQ ON MB SK AB BC
- <i>Diplotaxis erucoides</i> (L.) DC.	[rare] only Gaspé Pen, PQ
- <i>Diplotaxis muralis</i> (L.) DC. [sand-rocket]	NS PE NB PQ ON MB SK AB BC
- <i>Diplotaxis tenuifolia</i> (L.) DC. [wall-rocket]	NS NB PQ ON AB BC
- <i>Eruca vesicaria</i> (L.) Cav. ssp. <i>sativa</i> (Mill.) Thellung (= <i>E. sativa</i> Mill.) [garden-rocket]	[rare escape: PQ ON MB SK AB]
- <i>Erucastrum gallicum</i> (Willd.) O.E. Schulz [dog mustard]	NF NS PE NB PQ ON MB SK AB BC
- <i>Raphanus raphanistrum</i> L. [wild radish]	LB NF SM NS PE NB PQ ON (MB?) SK AB BC
- <i>Raphanus sativus</i> L. [radish]	NF NS PE NB PQ ON MB BC
- <i>Rapistrum perenne</i> (L.) All.	[rare] SK
- <i>Rapistrum rugosum</i> (L.) All.	[rare] PQ ON
- <i>Sinapis alba</i> L. [white mustard] (= <i>Brassica hirta</i> Moench)	YT NS PE NB PQ ON MB SK AB BC
- <i>Sinapis arvensis</i> L. [wild mustard] (= <i>Brassica kaber</i> (DC.) Wheeler)	NT-M YT LB NF SM NS PE NB PQ ON MB SK AB BC

CODE: Distribution: Northwest Territories, District of Mackenzie, NT-M; Yukon Territory, YT; Labrador, LB; Newfoundland, NF; Saint Pierre and Miquelon, SM; Prince Edward Island, PE; Nova Scotia, NS; New Brunswick, NB; Quebec, PQ; Ontario, ON; Manitoba, MB; Saskatchewan, SK; Alberta, AB; British Columbia, BC.

[REFERENCES: Crompton et al. 1988, Scoggan 1979]

Weed and crop species in United States: Distribution

- <i>Brassica fruticulosa</i> Cyr.	CA
- <i>Brassica juncea</i> (L.) Czern & Coss.	Almost every state
- <i>Brassica napus</i> L.	AK CA CT IL IN IA KY ME MD MA NY OH TN VT VA WV
- <i>Brassica nigra</i> (L.) Koch	Almost every state
- <i>Brassica oleracea</i> L.	CA CT IL IA KY MA NY OH RI TX VT
- <i>Brassica rapa</i> L. (= <i>B. campestris</i> L.)	Almost every state
- <i>Brassica tournefortii</i> Gouan	AZ CA TX
- <i>Cakile</i> spp.*	AK, Great Lakes, Atlantic and Pacific Coasts
- <i>Cakile maritima</i> Scop.	AL FL NJ
- <i>Coincya monensis</i> (L.) Greuter & Burdet	NC NY
- <i>Conringia orientalis</i> (L.) Andrzejowski	Almost every state
- <i>Crambe maritima</i> L.	OR
- <i>Diplotaxis muralis</i> (L.) DC.	AZ AR CA CT IL IN KS LA MA MI MN NE NJ NY NC OH OR PA SC SD TX UT WV
- <i>Diplotaxis tenuifolia</i> (L.) DC.	AZ CA CT DE IL IN KY MD MA MI NY OH OR PA TX VA
- <i>Eruca vesicaria</i> (L.) Cav. ssp. <i>sativa</i> (Mill.) Thellung (= <i>E. sativa</i> Mill.)	AZ CA CO CT IL IA KS MD MO MT NE NJ NM NY ND PA SD TX VT WA
- <i>Erucastrum gallicum</i> (Willd.) O.E. Schulz	CA FL ID IL IN KS KY LA ME MD MA MI MN MO MT NH NY ND PA SD TX VT WA WV WI WY
- <i>Hirschfeldia incana</i> (L.) Lagrèze-Fossat	CA NV
- <i>Raphanus raphanistrum</i> L.	Almost every state
- <i>Raphanus sativus</i> L.	Almost every state
- <i>Rapistrum rugosum</i> (L.) All.	CA LA MA NY PA TX
- <i>Sinapis alba</i> L. (= <i>Brassica hirta</i> Moench)	Almost every state
- <i>Sinapis arvensis</i> L. (= <i>Brassica kaber</i> L.)	Almost every state

CODE: Distribution: Alabama, AL; Alaska, AK; Arizona, AZ; Arkansas, AR; California, CA; Colorado, CO; Connecticut, CT; Delaware, DE; Florida, FL; Georgia, GA; Idaho, ID; Illinois, IL; Indiana, IN; Iowa, IA; Kansas, KS; Kentucky, KY; Louisiana, LA; Maine, ME; Maryland, MD; Massachusetts, MA; Michigan, MI; Minnesota, MN; Mississippi, MS; Missouri, MO; Montana, MT; Nebraska, NB; Nevada, NV; New Hampshire, NH; New Jersey, NJ; New Mexico, NM; New York, NY; North Carolina, NC; North Dakota, ND; Ohio, OH; Oklahoma, OK; Oregon, OR; Pennsylvania, PA; Rhode Island, RI; South Carolina, SC; South Dakota, SD; Tennessee, TN; Texas, TX; Utah, UT; Vermont, VT; Virginia, VA; Washington, WA; West Virginia, WV; Wisconsin, WI; Wyoming, WY.

[REFERENCES: A summary from all available floras, Al-Shehbaz 1985, Lemke & Worthington 1991]

*NOTE: The genus *Cakile* [North American component includes *C. edentula* (Bigel.) Hook., *C. constricta* Rodman, *C. geniculata* (Robinson) Millspaugh, *C. lanceolata* (Willd.) O.E. Schulz] is generally considered the only member of the Brassiceae with indigeneous species in the New World.

2. Morphological characters:

A. Hairiness (potential insect resistance):

Hairs on cotyledons (Gómez-Campo & Tortosa 1974, Warwick pers. observ.):

- <i>Boleum asperum</i>	(petiole & edge lamina)
- <i>Carrichtera annua</i>	(petiole)
- <i>Ceratocnemum rapistroides</i>	(petiole)
- <i>Crambe sventenii</i>	(petiole, all over lamina)
- <i>Otocarpus virgatus</i>	(petiole)
- <i>Psychine stylosa</i>	(petiole & edge lamina)
- <i>Sinapis alba</i>	(petiole and lamina)
- <i>Sinapis flexuosa</i>	(petiole and lamina)
- <i>Vella anrhemerica</i>	(petiole)
- <i>Vella pseudocytisus</i>	(petiole and lamina)
- <i>Vella spinosa</i>	(petiole)

Hairs very dense on juvenile leaves: (Gómez-Campo & Tortosa 1974)

- <i>Boleum asperum</i>	- <i>Crambe strigosa</i>
- <i>Brassica fruticulosa</i> ssp. <i>cossoniana</i>	- <i>Euzomodendron bourgaeum</i>
- <i>Brassica incana</i> (wild C)	- <i>Hirschfeldia incana</i>
- <i>Brassica villosa</i> (wild C)	- <i>Sinapis alba</i>
- <i>Coincya monensis</i>	- <i>Sinapis flexuosa</i>
- <i>Coincya rupestris</i>	- <i>Sinapis pubescens</i>
- <i>Crambe arborea</i>	- <i>Vella pseudocytisus</i>
- <i>Crambe orientalis</i>	- <i>Vella spinosa</i>

Hairs on fruit: (Warwick pers. observ.)

- <i>Boleum asperum</i>
- <i>Carrichtera annua</i>
- <i>Ceratocnemum rapistroides</i>
- <i>Cordylocarpus muricatus</i>
- <i>Enarthrocarpus arcuatus</i>
- <i>Guiraoa arvensis</i>
- <i>Muricaria prostrata</i>
- <i>Otocarpus virgatum</i>
- <i>Psychine stylosa</i>
- <i>Rapistrum rugosum</i>
- <i>Sinapis alba</i>
- <i>Sinapis flexuosa</i>
- <i>Trachystoma aphanoneurum</i> (antrose hair)
- <i>Trachystoma ballii</i> (retrosc hair)
- <i>Trachystoma labasii</i> (antrose hair)
- <i>Vella mairei</i>
- <i>Vella pseudocytisus</i>

B. Leaf thickness/waxiness: [drought & insect tolerance]:

- <i>Brassica cretica</i>
- <i>Brassica oleracea</i> (Stoner 1990)
- <i>Diplotaxis harra</i>
- <i>Moricandia</i> spp.
- <i>Rytidocarpus</i>

C. Petal colour (Yellow with the following exceptions):**White:**

- Brassica "alboglabra"*
- Brassica cretica* ssp. *cretica*
- Brassica insularis*
- Diplotaxis erucoides*
- Erucastrum leucanthum*

Purple/White:

- subtribe Cakilinae (*Cakile*, *Didesmus*, *Erucaria*, *Reboudia*)

Purple:

- Eruca setulosa*
- Moricandia* spp. & *Rytidocarpus*
- subtribe Zillinae (*Foleyola*, *Fortuynia*, *Physorrhynchus*, *Schouwia*, *Zilla*)

Apetalous flower:

[NOTE: Problem with photosynthetic efficiency of rapeseed due to dense floral canopy which can reflect or absorb as much as 60% of the solar radiation (Rao et al. 1991).]

- Conringia orientalis* [petals very small and occasionally absent] (Rich 1991)

D. Resistance to pod shattering:

- | | |
|--------------------------------|--------------------------|
| - <i>Brassica juncea</i> | (Prakash & Chopra 1988a) |
| - <i>Brassica macrocarpa</i> * | (Mithen & Herron 1991) |
| - <i>Brassica hilarionis</i> * | (Mithen & Herron 1991) |
| - <i>Brassica tournefortii</i> | (Salisbury 1989) |
| - <i>Conringia orientalis</i> | (Salisbury 1989) |
| - <i>Enarthrocarpus</i> spp. | (Warwick pers. observ.) |
| - <i>Hirschfeldia incana</i> | (Salisbury 1989) |
| - <i>Raphanus</i> spp. | (Agnihotri et al. 1991) |

*: partial resistance

E. Yellow seed: (Sobrino-Vesperinas et al. 1991):

- Brassica carinata*
- Brassica juncea*
- Brassica rapa*
- Sinapis alba*

F. Growth form:

Annual, winter annuals, biennials, perennials, woody shrubs

Sub-trees:

Canary Island *Crambe* spp. (eg. *C. arborea* & *C. strigosa*)

Burial of fruit underground (geogarpy):

- Morisia monanthos*
- Raffenaldia primuloides*

3. Chemical Traits:

A. Fatty Acids: (Reviewed by Kumar & Tsunoda (1980) for tribe):

There is variation in both the amount of oil and fatty acid composition.

Additional surveys of wild species:

Vioque et al. (1990) - 4 wild *Brassica* spp.

Yaniv et al. (1991) - 11 wild taxa in tribe

High erucic acid (22:1) (>45-50%):

-*Brassica cretica* (Yaniv et al. 1991)

-*Crambe abyssinica* and *C. hispanica* (Yaniv et al. 1991)

-*Eruca vesicaria* (includes *E. sativa*) (Yaniv et al. 1991)

-*Sinapis alba* (Persson 1986, Yaniv et al. 1991)

B. Glucosinolates: (Mustard oil glucosides)

Characteristic odours and flavours of crucifers are due to the glucosinolate hydrolysis products. These have long been of toxicological interest (Daxenbichler et al. 1991), as certain glucosinolates and their degradation products have shown adverse effects on animals. More recently there is an interest in their pharmacological role in the prevention of disease and in their use as insect attractants/repellants, etc.. Greater than 60 glucosinolates are present in the Cruciferae, and many of these are unique to certain species and genera.

Surveys of glucosinolates in wild members of the tribe:

Al-Shenhbaz & Al-Shammery (1987) - 13 species [total of 51 cruciferous species]

Daxenbichler et al. (1991) - 64 species [total of 259 cruciferous species]

El-Menshawhi et al. (1980) - 3 species of *Zilla*

Horn & Vaughan (1983) - 14 species of *Brassica*

Mithen et al. (1987a) - 9 species of the wild *Brassica oleracea* complex

Rodman (1974, 1976) - 13 species of *Cakile*

C. Aromatic Phenolic Choline Esters (such as sinapine):

Bouchereau et al. (1991) surveyed 23 taxa in Tribe and found both specific and generic differences.

4. Photosynthesis (All C_3 plants with following exception):

C_3 - C_4 Intermediate:

-*Moricandia* spp. [*M. arvensis*, *M. nitens*, *M. sinaica*, *M. spinosa*, and *M. suffruticosa*]

(Bauwe 1983, Hylton et al. 1988)

5. Cytoplasmic male sterility:

-*Brassica oxyrrhina* (Prakash & Chopra 1988b)

-*Brassica tournefortii* (Pradhan et al. 1991)

-*Diplotaxis harra* (Klimaszewska & Keller 1988)

-*Diplotaxis muralis* (Hinata & Konno 1979)

-*Hirschfeldia incana* (Horovitz & Galil 1972)

-*Raphanus sativus* (Ogura 1968)

6. Soil adaptation:

A. Tolerance to saline soils:

- (succulent annuals of coastal strand habitats)
 -*Cakile maritima** [North American component includes *C. edentula*]
 -*Crambe maritima*
 -*Raphanus raphanistrum* ssp. *maritimus* [=*Raphanus maritimus*]

NOTE: Unique ecological specialization for these three groups is unknown elsewhere in the Cruciferae (Al-Shehbaz 1985).

*NOTE: Habitat for all members of genus *Cakile* with exception of *C. arabica*

B. Tolerance to calcareous (limestone) soils:

- Euzomodendron bourgaeum*
 -*Moricandia moricandiodes*

7. Cold tolerance:

Mountainous habitats (elevations > 2000-2500 m or above snow line)

- Brassica nivalis* (Mt. Olympus, Greece)
 -*Brassica jordanoffii* (Mt Pirin Planina, s.w. Bulgaria)
 -*Coincya richeri* (S.W. Alps in France & Italy- occurs above snowline 1750-2500m (Leadlay & Heywood 1990)

-*Erucastrum abyssinicum* & *E. pachypodum*, Ethiopian Highlands (> 3000m) (Al-Shehbaz 1985)

8. Drought tolerance (desert endemics):

A. Annuals:

- | | |
|---|--------------------------|
| - <i>Brassica juncea</i> | (Downey & Röbbelen 1989) |
| - <i>Brassica tournefortii</i> | (Salisbury 1989) |
| - <i>Carrichtera annua</i> | (Boaz et al. 1990) |
| - <i>Diploaxis acris</i> & <i>D. harra</i> | (Boaz et al. 1990) |
| - <i>Enarthrocarpus strangulatus</i> | (Boaz et al. 1990) |
| - <i>Eruca sativa</i> | (Sun et al. 1991) |
| - <i>Erucaria boveana</i> | (Boaz et al. 1990) |
| - <i>Erucaria microcarpa</i> & <i>E. uncata</i> | (Boaz et al. 1990) |
| - <i>Moricandia</i> spp. | (McVetty et al. 1989) |
| - <i>Pseuderucaria clavata</i> | (Boaz et al. 1990) |
| - <i>Savignya parviflora</i> | (Boaz et al. 1990) |
| - <i>Schouwia purpurea</i> | (Boaz et al. 1990) |
| - <i>Sinapis alba</i> | (Downey & Röbbelen 1989) |

B. Desert shrubs:

- subtribe Zillinae
 (*Foleyola*, *Fortuynia*, *Physorrhynchus*, *Zilla*)

9. Herbicide resistance:

A. Atrazine resistance:

- Brassica rapa* (Quebec) (Warwick 1991, Warwick & Black 1993b)
 -*Sinapis arvensis* (Ontario) (Warwick 1991)

B. Phenoxy resistance:

- Sinapis arvensis* (Manitoba) (Morrison, pers. comm.)

10. Disease resistance:

A. *Albugo candida* -- White rust

Resistance to *A. candida* races/isolates has been found at both the *Brassica* species and cultivar levels.

- Brassica napus*: Canadian cultivars highly resistant to western Canadian isolates (Liu & Rimmer 1991)
- Brassica carinata*: resistance common, *B. rapa*, *B. juncea*, *B. nigra*: resistance more limited in latter three (Liu & Rimmer 1991, Gulati et al. 1991, Kolte et al. 1991)
- Raphanus sativus* (Williams & Pound 1963, Kolte et al. 1991)

B. *Alternaria brassicae* -- Blackspot/ *Alternaria* blight

- Brassica spinescens* (Agnihotri et al. 1991)
- Eruca sativa*: highly resistant to a Canadian isolate (Conn & Tewari 1986, Tewari 1991)
- Sinapis alba* (Kolte 1985, Brun et al. 1987, Sharma & Singh 1992)

[NOTE: Highest resistance in crucifers found in *Camelina sativa* (False flax) and *Capsella bursa-pastoris* (Shepherd's purse) (Conn et al. 1988, reviewed in Tewari 1991)]

C. *Leptosphaeria maculans* [=Phoma lingam] -- Blackleg

- Brassica carinata*, *B. juncea*, *B. nigra* (Rimmer & van den Berg 1992)
- Brassica insularis*, *B. atlantica*, *B. macrocarpa*: Members of wild *B. oleracea* complex] (Mithen et al. 1987b, Mithen & Herron 1991, Mithen & Magrath 1992)
- Weedy species of *Carrichtera*, *Diplotaxis*, *Hirschfeldia*, *Raphanus*, *Rapistrum* and *Sinapis* (Salisbury 1987)

D. *Peronospora parasitica* -- Downy mildew

- wild *Brassica oleracea* (Greenhalgh & Mitchell 1976)

E. *Plasmodiophora brassicae* -- Clubroot

- Raphanus* spp. (Crute et al. 1980, Long et al. 1992)

F. Virus Resistance(Immunity):

Turnip mosaic virus (common strain TuMV-S1)

- Raphanus raphanistrum* (Stobbs & Stirling 1990; surveyed 6 weedy species in the tribe]

G. *Sclerotinia sclerotiorum* -- Sclerotinia stem rot

- [NOTE: occurs worldwide, infects at least 225 genera of host plants, including weeds (Purdy 1979) can cause serious losses in yield of canola due to lodging and premature shattering of seedpods]
- no known source of resistance

11. Insect resistance:

A. Flea beetle [*Phyllotreta cruciferae* and *P. striolata*]

Flea beetles are the most important insect pests of oilseed crucifer crops in the Prairies (Canada) and Northern Great Plains (USA). Oilseed rape is most vulnerable in the spring when adult flea beetles attack newly emerged seedlings producing pits on leaves and cotyledons, that when severe can cause seedling mortality.

- Brassica incana* (Bodnaryk in prep.)
- Brassica villosa* (Bodnaryk in prep.)
- Crambe abyssinica* (Anderson et al. 1992)
- Sinapis alba* (Bodnaryk & Lamb 1991, Lamb 1980)

B. Lepidopterous pests --

- Brassica oleracea* - Glossiness (Stoner 1990)

C. Cabbage aphid [*Brevicoryne brassicae*]

- Sinapis alba* (Thompson 1963)

12. Nematode resistance:

Beet cyst nematode (*Heterodera schachtii*)

- Raphanus sativus* into *Brassica napus*
(Thierfelder et al. 1991, Lelivelt & Krens 1992)
- Sinapis alba* into *Brassica napus* (Lelivelt et al. 1993)
- Sinapis alba* (Thierfelder et al. 1991)

13. New crops:

Source for vegetables, fodder, condiments and oil

- "Brassicoraphanus": (raparadish) (Lange et al. 1989)
- Crambe abyssinica*: (Erickson & Bassin 1990)
Insect tolerant & high levels of erucic acid (industrial oil)
- Eruca sativa*: (Sun et al. 1991)
Drought tolerant & high levels of erucic acid (industrial oil)
- Sinapis alba*: (Downey & Röbblen 1989)
Condiment mustard - c. 57,000 ha grown annually in Canada
Drought tolerance and tolerance to high temperatures, insect tolerance

REFERENCES

- Agnihotri, A., K.R. Shivanni, M.S. Lakshmikumaran & V. Jagannathan (1991) Micropropagation and DNA analysis of wide hybrids of cultivated *Brassica*. GCIRC 1991 Congress, Abst. pp: 151.
- Al-Shehbaz, I.A. (1984) The tribes of Cruciferae (Brassicaceae) in the southeastern United States. *J. Arnold Arboretum* 65: 343-373.
- Al-Shehbaz, I.A. (1985) The genera of Brassiceae (Cruciferae: Brassicaceae) in the southwestern United States. *J. Arnold Arboretum* 66: 279-351.
- Al-Shehbaz, I.A. & K.I. Al-Shammmary (1987) Distribution and chemotaxonomic significance of glucosinolates in certain Middle-Eastern Cruciferae. *Biochem. System. & Ecol.* 15: 559-569.
- Anderson, M.D., C. Peng & M.J. Weiss (1992) *Crambe*, *Crambe abyssinica* Hochst., as a flea beetle resistant crop (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* 85: 594-600.
- Arnason, T., R.J. Hebda & T. Johns (1981) Use of plants for food and medicine by Native Peoples of eastern Canada. *Can. J. Bot.* 59: 2189-2325.
- Bauwe, H. (1983) Comparative phylogenetic age of C₃-C₄ intermediate species of *Moricandia* determined by isoelectric focusing and amino acid composition of small subunit of ribulose 1,5-bisphosphate carboxylase/oxygenase. *Photosynthetica* 17: 442-449.
- Bing, D.J., R.K. Downey & G.F.W. Rakow (1991) Potential of gene transfer among oilseed *Brassica* and their weedy relatives. GCIRC 1991 Congress, pp: 1022-1027.
- Boaz, M, U. Plitmann & C.C. Heyn (1990) The ecogeographic distribution of breeding systems in the Cruciferae (Brassicaceae) of Israel. *Israel J. Bot.* 39: 31-42.
- Bodnaryk R.P. & R.J. Lamb (1991) Mechanisms of resistance to the flea beetle, *Phyllotreta cruciferae* (Goeze), in yellow mustard seedlings, *Sinapis alba* L.. *Can. J. Plant Sci.* 71: 13-20.
- Bouchereau A., J. Hamelin, I. Lamour, M. Renard & F. Larher (1991) Distribution of sinapine and related compounds in seeds of *Brassica* and allied genera. *Phytochemistry* 30: 1873-1881.
- Brun H., J. Pleiss & M. Renard (1987) Resistance of some crucifers to *Alternaria brassicae* (Berk.) Sacc.. 7th International Rapeseed Congress. Poznan 11-14 May 1987, pp: 1222-1227.
- Conn, K.L. & J.P. Tewari (1986) Hypersensitive reaction induced by *Alternaria brassicae* in *Eruca sativa*, an oil yielding crucifer. *Can. J. Plant Path.* 8: 348.
- Conn, K.L., J.P. Tewari & J.S. Dahiya (1988) Resistance to *Alternaria brassicae* and phytoalexin-elicitation in rapeseed and other crucifers. *Plant Sci.* 56: 21-25.
- Crompton, C.W., J. McNeil, A.E. Stahevitch & W.A. Wojtas (1988) Preliminary inventory of Canadian weeds. Research Branch, Agriculture Canada. Tech. Bull. 1988-9E, 291 pp.

- Crute I.A., A.R. Gray, P. Crisp & S.T. Buczacki (1980) Variation in *Plasmodiophora brassicae* and resistance to clubroot disease in *Brassicaceae* and allied crops. Pl. Breeding Abstracts 50: 91-104.
- Daxenbichler, M.E., G.F. Spencer, D.G. Carlson, G.B. Rose, A.M. Brinker & R.G. Powell (1991) Glucosinolate composition of seeds from 297 species of wild plants. Phytochemistry 30: 2623-2638.
- Downey, R.K. & G. Röbbelen (1989) *Brassica* species In: Oil Crops of the World. Edited by G. Röbbelen, R.K. Downey & A. Ashri. McGraw-Hill, New York. pp: 339-362.
- El-Menshaw, B., M. Karawya, G. Wassel, J. Reisch & A. Kjaer (1980) Glucosinolates in the genus *Zilla* (*Brassicaceae*). J. Natural Products 43: 534-536.
- Erickson, D.B. & P. Bassin (1990) Rapeseed and Crambe: alternative crops with potential industrial uses. Bull. Kansas Agric. Expt. Stn. (Manhattan) No. 656: 36 pp.
- Gómez-Campo, C. (1980) Morphology and morphotaxonomy of the Tribe *Brassicaceae*. In: *Brassica* crops and wild allies. Edited by S. Tsunoda, K. Hinata & C. Gómez-Campo. Japan Science Societies Press, Tokyo. pp: 3-31.
- Gómez-Campo C. & M.E. Tortosa (1974) The taxonomic and evolutionary significance of some juvenile characters in the *Brassicaceae*. Bot. J. Linn. Soc. 69: 105-124.
- Greenhalgh J.G. & N.D. Mitchell (1976) The involvement of flavour volatiles in the resistance of downy mildew of wild and cultivated forms of *Brassica oleracea*. New Phytol. 77: 391-398.
- Gulati, S.C., N.S. Varma, N. Mani & R. Raman (1991) Resistance to white rust (*Albugo candida*) in Indian mustard. GCIRC 1991 Congress, pp: 256-261.
- Hedge, I.C. (1976) A systematic and geographical survey of the Old World *Cruciferae*. In: The biology and chemistry of the *Cruciferae*. Edited by J.G. Vaughn, A.J. MacLeod & B.M.G. Jones. Academic Press, London. pp: 1-45.
- Hinata K. & N. Konno (1979) Studies on a male sterile strain having the *Brassica campestris* nucleus and the *Diplotaxis muralis* cytoplasm I. On the breeding procedure and some characteristics of the male sterile strain. Jap. J. Breed. 29: 305-311.
- Horn, P.J. & J.G. Vaughn (1983) Seed glucosinolates of fourteen wild *Brassica* species. Phytochemistry 22: 465-470.
- Horovitz, A. & J. Galil (1972) Gynodioecism in east Mediterranean *Hirschfeldia incana*, *Cruciferae*. Bot. Gaz. 133: 127-131.
- Hylton, C.M., S. Rawsthorne, A.M. Smith, D.A. Jones & H.W. Woolhouse (1988) Glycine decarboxylase is confined to the bundle-sheath cells of leaves of C_3 - C_4 intermediate species. Planta 175: 452-459.
- Jacobson, H.A., J.B. Petersen & D.E. Putnam (1988) Evidence of pre-Columbian *Brassica* in the northeastern U.S.. Rhodora 90: 355-362.
- Kerlan, M.C., A.M. Chevre, F. Eber, J. Botterman & W. de Greef (1991) Risk assessment of gene transfer from transgenic rapeseed to wild species in optimal conditions. GCIRC 1991 Congress, pp: 1028-1033.

- Klimaszewska, K. & W.A. Keller (1988) Regeneration and characterization of somatic hybrids between *Brassica napus* and *Diplotaxis harra*. Pl. Sci. 58: 211-222.
- Kolte, S.J. (1985) Diseases of annual edible oilseed crops. Vol. 2-3. CRC Press, Boca Raton, Fla.
- Kolte, S.J., D.K. Bordoloi & R.P. Awasthi (1991) The search for resistance to major diseases of rapeseed and mustard in India. GCIRC 1991 Congress, pp: 219-225.
- Kumar, P.R. & S. Tsunoda (1980) Variation in oil content of fatty acid composition among seeds from the Cruciferae. In: *Brassica crops and wild allies*. Edited by S. Tsunoda, K. Hinata & C. Gómez-Campo. Japan Scientific Societies Press, Tokyo. pp: 235-252.
- Lange, W., H. Toxopeus, J.H. Lubberts, O. Dolstra & J.L. Harrewijn (1989) The development of raparadish *Brassicoraphanus* 2n equals 38, a new crop in Agriculture. Euphytica 40: 1-14.
- Lamb, R.J. (1980) Hairs protect pods of mustard (*Brassica hirta* 'Gisilba') from flea beetle feeding damage. Can. J. Plant Sci. 60: 1439-1440.
- Leadlay, E.A. & V.H. Heywood (1990) The biology and systematics of the genus *Coincya* Porta & Rigo ex. Rouy (Cruciferae). Bot. J. Linn. Soc. 102: 313-398.
- Lefol, E., V. Danielou & D. Darmency (1991) Escape of engineered genes from rapeseed to wild Brassiceae. Brighton Crop Protection Conference - Weeds-1991, pp: 1049-1056.
- Lelivelt, C.L. & F.A. Krens (1992) Transfer of resistance to the beet cyst nematode *Heterodera schachtii* Schm. into the *Brassica napus* L. gene pool through intergeneric somatic hybridization with *Raphanus sativus* L.. Theor. Appl. Genet. 83: 887-894.
- Lelivelt, C.L.C., E.H.M. Leunissen, H.J. Frederiks, J.P.F.G. Helsper & F.A. Krens (1993) Transfer of resistance of the beet cyst nematode (*Heterodera schachtii* Schm.) from *Sinapis alba* L. (white mustard) to the *Brassica napus* L. gene pool by means of sexual and somatic hybridization. Theor. Appl. Genet. 85: 688-696.
- Lemke D.E. & R.D. Worthington (1991) *Brassica* and *Rapistrum* (Brassicaceae) in Texas. The Southwestern Naturalist 36: 194-197.
- Liu, Q. & S. R. Rimmer (1991) Inheritance of resistance in *Brassica napus* to an Ethiopian isolate of *Albugo candida* from *Brassica carinata*. Can J. Plant Pathol. 13: 197-201.
- Long, M.L., G.M. Xing, H. Okubo & K. Fujieda (1992) Cross compatibility between *Brassicoraphanus* (*Brassica oleracea* X *Raphanus sativus*) and cruciferous crops, and rescuing the hybrid embryos through ovary and embryo culture. J. Fac. Agr. Kyushu Univ. 37: 29-39.
- McVetty, P.B.E., R.B. Austin & C.L. Morgan (1989) A comparison of the growth, photosynthesis, stomatal conductance and water use efficiency of *Moricandia* and *Brassica* species. Annals of Botany 64: 87-94.
- Mithen, R.F., B.G. Lewis, R.K. Heaney & G.R. Fenwick (1987a) Glucosinolates of wild and cultivated *Brassica* species. Phytochemistry 26: 1969-1973.

- Mithen, R.F., B.G. Lewis, R.K. Heaney & G.R. Fenwick (1987b) Resistance of leaves of *Brassica* species to *Leptosphaeria maculans*. Trans. Br. Mycol. Soc. 88: 525-531.
- Mithen, R.F. & C. Herron (1991) Transfer of disease resistance to oilseed rape from wild *Brassica* species. GCIRC 1991 Congress, pp: 244-249.
- Mithen, R.F. & R. Magrath (1992) Glucosinolates and resistance to *Leptosphaeria maculans* in wild and cultivated *Brassica* spp.. Plant Breeding 108: 60-68.
- Ogura, H. (1968) Studies on the new male-sterility in Japanese radish, with special reference to the utilization of this sterility towards the practical raising of hybrid seeds. Mem. Fac. Agric. Kagoshima Univ. 6: 39-78.
- Panetsos, C. & H.G. Baker (1968) The origin of variation in "wild" *Raphanus sativus* (Cruciferae) in California. Genetica 38: 243-274.
- Persson, C. (1986) High erucic oil from white mustard (*Sinapis alba* L.) for technical use. Cruciferae Newsletter 11: 134.
- Pradhan, A.K., A. Mukopadhyay & D. Pental (1991) Identification of the putative cytoplasmic donor of a cms system in *Brassica juncea*. Plant Breeding 106: 204-208.
- Prakash, S. & V.L. Chopra (1988a) Introgression of resistance to pod shatter in *Brassica napus* from *Brassica juncea* through non-homologous recombination. Plant Breeding 101: 167-168.
- Prakash, S. & V.L. Chopra (1988b) Synthesis of alloplasmic *Brassica campestris* and induction of cytoplasmic male sterility. Plant Breeding 101: 253-255.
- Purdy, L.H. (1979) *Sclerotinia sclerotiorum*: history, diseases and symptomology, host range, geographic distribution and impact. Phytopathology 69: 875-880.
- Rao, M.S.S., N.J. Mendham & G.C. Buzza (1991) Effect of the apetalous flower character on radiation distribution in the crop canopy yield and its components in oilseed rape *Brassica napus*. J. Agric. Sci. 117: 189-196.
- Rich, T.C.G. (1991) Crucifers of Great Britain and Ireland. Botanical Society of the British Isles. Handbook No. 6, London, U.K.. 336 pp.
- Rimmer S.R. and C.G.J. van den Berg (1992) Resistance of oilseed *Brassica* spp. to blackleg caused by *Leptosphaeria maculans*. Can. J. Pathol. 14: 56-66.
- Rodman, J.E. (1974) Systematics and evolution of the genus *Cakile* (Cruciferae). Contr. Gary Herb. Harv. Univ. 205: 3-146.
- Rodman, J.E. (1976) Differentiation and migration of *Cakile* (Cruciferae): seed glucosinolate evidence. Syst. Bot. 1: 137-148.
- Salisbury, P.A. (1987) Blackleg resistance in weedy crucifers. Cruciferae Newsletter 12: 90.
- Salisbury, P.A. (1989) Potential utilization of wild crucifer germplasm in oilseed *Brassica* breeding. Proceedings of the 7th Australian Rapeseed Agronomists and Breeders Workshop, pp: 51-53.
- Schulz, O.E. (1919) IV. 105 Cruciferae-Brassicaceae. Part 1. Subtribes Brassicinae and Raphaninae. In: Das Pflanzenreich. Edited by A. Engler, Heft 68-70, Wilhelm Engelmann, Leipzig. pp: 1-290.

- Schulz, O.E. (1923) IV. 105. Cruciferae-Brassicaceae. Part II. Subtribes Cakilinae, Zillinae, Vellinae, Savignyinae and Moricandiinae. In: Das Pflanzenreich. Edited by A. Engler, Heft 82-85, Wilhelm Engelmann, Leipzig. pp: 1-100.
- Schulz O.E. (1936) In: Die Natürlichen Pflanzenfamilien, 2nd Ed. Edited by A. Engler & K. Prantl, Band 17-b, Wilhelm Engelmann, Leipzig. pp: 227-658.
- Scoggan, H.G. (1979) The flora of Canada. Vol. 3, Cruciferae. Nat. Mus. Nat. Sci. (Ottawa), Publ. Bot. 7. National Museums of Canada, Ottawa, Ontario.
- Sharma T.R. & B.M. Singh (1992) Transfer of resistance to *Alternaria brassicae* in *Brassica juncea* through interspecific hybridization among *Brassica*. J. Genet. & Breed. 46: 373-378.
- Sjödin, C. (1992) Brassicaceae, a plant family well suited for modern biotechnology. Acta. Agric. Scand., Sect. B. Soil and Plant Sci. 42: 197-207.
- Sobrinho-Vesperinas, E., M.C. Fernandez & F. Monroy (1991) Characteristics of a yellow-seed line belonging to *Brassica napus* L. GCIRC 1991 Congress, pp: 203-210.
- Stobbs, L.W. & A. Stirling (1990) Susceptibility of Ontario weed species to turnip mosaic virus. Can. J. Plant Pathol. 12: 255-262.
- Stoner, K. (1990) Glossy leaf wax and plant resistance to insects in *Brassica oleracea* under natural infestation. Environmental Entomol. 19: 730-739.
- Sun, W.C., Q.Y. Pan, X.H. An & Y.P. Yang (1991) *Brassica* and *Brassica*-related oilseed crops in Gansu, China. GCIRC 1991 Congress: 1130-1135.
- Tewari, J.P. (1991) Current understanding of resistance to *Alternaria brassicae* in Cruciferae. GCIRC 1991 Congress, pp: 471-476.
- Thierfelder, A., E. Hackenberg, K. Nichterlein & W. Friedt (1991) Development of nematode-resistant rapeseed genotypes via interspecific hybridization. GCIRC 1991 Congress, pp: 269-273.
- Thompson, K.F. (1963) Resistance to the cabbage aphid (*Brevicoryne brassicae*) in brassica plants. Nature 198: 209.
- Vioque, J., J. Pastor & E. Vioque (1990) Analysis of the fatty acids and sterols of the oils of four *Brassica* species. Lagasalia 16: 95-103.
- Warwick, S.I. (1991) Herbicide resistance in weedy plants: physiology and population biology. Annu. Rev. Ecol. Syst. 22: 95-114.
- Warwick, S.I. & L.D. Black (1991) Molecular systematics of *Brassica* and allied genera (Subtribe Brassicinae, Brassicaceae) - chloroplast genome and cytodeme congruence. Theor. Appl. Genet. 82: 81-92.
- Warwick, S.I. & L.D. Black (1993a) Molecular relationships in subtribe Brassicinae (Cruciferae, tribe Brassicaceae). Can. J. Bot. 71: 906-918.
- Warwick, S.I. & L.D. Black (1993b) Electrophoretic variation in triazine-resistant and susceptible populations of the allogamous weed *Brassica rapa*. Weed Res. 33: 105-114.
- Williams, P.H. & G.S. Pound (1963) Nature and inheritance of resistance to *Albugo candida* in radish. Phytopathology 53: 1150-1154.
- Yaniv, Z, Y. Elber, M. Zur & D. Schafferman (1991) Differences in fatty acid composition of oils of wild cruciferae seed. Phytochemistry 30: 841-843.

CANADIAN AGRICULTURE LIBRARY



BIBLIOTHEQUE CANADIENNE DE L'AGRICULTURE

3 9073 00107321 4

